

Original Research Article

HPLC-UV-MS/MS Profiling of Phenolics from *Euphorbia nicaeensis* (All.) Leaf and Stem and Its Antioxidant and Anti-Protein Denaturation Activities

Khawla Bouaouda¹, Chaimae Elagdi², Naoufal El Hachlafi³, Karima Mohtadi¹, Mohammed Hsaine², Anass Kettani¹, Rachid Flouchi³, Khang Wen Goh⁴, Abdelhakim Bouyahya⁵, Hanae Naceiri Mrabti⁶, Rachid Saile¹, Hassan Taki^{1*}

Article History

Received: 19 March 2023;

Received in Revised Form:

21 April 2023;

Accepted: 10 May 2023;

Available Online: 05 June 2023

¹Laboratory of Biology and Health, University Hassan II of Casablanca, Faculty of Sciences Ben M'Sik, Casablanca, Morocco, P.O.Box 7955; khawla.bouaouda-etu@etu.univh2c.ma (KB); karima.mohtadi@univh2c.ma (KM); anass.kettani@univh2c.ma (AK); rachid.saile@univh2c.ma (RS)

²Laboratory of Ecology and Environment, University Hassan II of Casablanca, Faculty of Sciences Ben M'Sik, Casablanca, Morocco, P.O.Box 7955; chaimaa.elagdi-etu@etu.univh2c.ma (CE); mohammed.hsaine@univh2c.ma (MH)

³Laboratory of Microbial Biotechnology and Bioactive Molecules, Sciences and Technologies Faculty, Sidi Mohamed Ben Abdellah University, P.O. Box 2202, Imouzzar Road, Fez, Morocco; naoufal.elhachlafi@usmba.ac.ma (NEH), racinf@gmail.com (RF)

⁴Faculty of Data Science and Information Technology, INTI International University, 71800 Nilai, Malaysia; khangwen.goh@newinti.edu.my (KWG)

⁵Laboratory of Human Pathologies Biology, Department of Biology, Faculty of Sciences, Mohammed V University in Rabat, Rabat 10106, Morocco; a.bouyahya@um5r.ac.ma (AB)

⁶High Institute of Nursing Professions and Health Techniques Casablanca, Morocco; naceiri.mrabti.hanae@gmail.com (HNM)

*Corresponding author: Hassan Taki, University Hassan II of Casablanca, Faculty of Sciences Ben M'Sik, Casablanca, Morocco; hassan.taki@univh2c.ma (HT)

Abstract: This work focuses on the leaves and stems of *Euphorbia nicaeensis* All. to confirm its historical use by the Moroccan population. The phytochemical profile of the plant by HPLC-UV-MS/MS was identified for the first time, and the biological activities of each plant organ was evaluated separately by maceration and ultrasonic-assisted extraction. The evaluation of antioxidant activity based on DPPH assay and hydrogen peroxide scavenging assay showed that leaves could be used as a natural source of antioxidants as they provide a potent antioxidant effect (DPPH IC₅₀ = 23.47±0.62 µg/ml), (H₂O₂ IC₅₀ = 110.27 ± 3.59 µg/ml) compared to stems. These results were proven by HPLC-UV-MS/MS analysis and revealed that *E. nicaeensis* leaves are richer in phenolic compounds, especially quercetin and derivatives, known for their antioxidant properties. In contrast, the stems could be considered a potential anti-inflammatory agent considering their solid anti-inflammatory activity. The

most potent effects were obtained at a concentration of 2 mg/ml, which induced 83.98% and 82.04% inhibition against bovine albumin and egg albumin denaturation, respectively, compared to the control. In addition, the stem phytochemical profile indicated the presence of some compounds with anti-inflammatory effects, such as fargesin and nuciferine. Likewise, the findings showed that ultrasound-assisted extraction was more effective than maceration.

Keywords: *Euphorbia nicaeensis* All., Maceration, Ultrasound extraction, Antioxidant activity, Anti-inflammatory activity, HPLC-UV-MS/MS

1. Introduction

Aromatic and medicinal plants have long been regarded as a vital source of therapeutics and curative remedies due to the presence of bioactive compounds and phytochemical constituents. Medicinal herbs have proven to be the basis of traditional medicine worldwide. Curing and healing human diseases has always been related to using entire plants or parts of them in remedy preparation ^[1].

Moroccan population has traditionally employed aromatic and medicinal plants; over time, people have gained knowledge of old cosmetics and pharmacopoeia. Oral communication is still used to pass on essential information about these activities from generation to generation ^[2]. However, several medicinal plants are currently underexploited in Morocco despite their ancestral use in treating human diseases.

Euphorbia nicaeensis All. is one of these plants on which this investigation focuses. It is a species of Euphorbiaceae family, a perennial spurge herbaceous plant that prefers calcareous soils. It grows in sunny and dry places and is distributed in the Mediterranean and central Europe with a strong morphological variability in leaves and bracts ^[3].

E. nicaeensis is an abundant species in the rocky lawns of Morocco ^[4], latex is the most traditionally used part of the plant, and it has been used in the past to attack warts and erase dead flesh. The population was aware of the vesicant properties of latex on the skin, eyes, and mucous membranes ^[5]. There are also other uses of leaves and stems against bacterial infections, hepatitis, tuberculosis, typhoid, and other diseases ^[6–8].

Numerous studies have proven the traditional use of *E. nicaeensis* and its anthelmintic, antifungal, and anticancer activities due to the jatrophone diterpenoid isolated from the root extracts and latex ^[9–11]. Previous studies have also evaluated some biological activities of *E. nicaeensis* aerial parts and detected its anti-inflammatory properties ^[12,13]. This study aims to assess the phytochemical profile of *Euphorbia nicaeensis* by HPLC-UV-MS/MS analysis for the first time and to evaluate the anti-inflammatory properties and antioxidant activity of leaves and stems. Therefore, two extraction methods were compared, probe ultrasonic-assisted extraction and maceration.

2. Materials and Methods

2.1. Plant material

The plant was harvested from the Ifrane region (Figure 1). The Lambert coordinates: 33° 32' 44.4" N, 5° 19' 17.89" W. and the plant have undergone drying after separation of the leaves.



Figure 1. Photo of the species *Euphorbia nicaeensis* All.

2.2. Extraction procedure

2.2.1. Maceration

An amount of 2.5 g of ground plant powder (leaves and stems) was extracted by 50 ml (methanol - water) (80:20) v/v. The maceration extracts were kept under stirring for 24 hours, then filtered under a Buchner funnel and concentrated by rotary evaporation. The residues were collected and stored until further analysis.

2.2.2. Probe ultrasonic-assisted extraction (PUAE)

Probe ultrasonic-assisted extraction (PUAE) is one of the main extraction methods used for plant materials. Sonication causes cavitation and implosion, which causes cell-wall rupture and increases the number of disturbed cells. When disturbed, the solvent penetrates the cell, and the intracellular plant material is absorbed into the solvent ^[14].

The extraction was done using an ultrasonic probe system (Bioblock scientific, Vibra Cell 75042). The probe was submerged 1.5 cm under the surface of the mixture (methanol-water) (80:20) v/v and 2.5 g of plant powder. The extraction was performed at the maximum power settings of the transducer (100%, 400 W), at 24 kHz, for 15 min, with a 20 s pulse. Following extraction, the extract was filtered and concentrated by rotary evaporation until dryness and was stored for further use.

2.3. Phytochemical screening

Phytochemical screening of *Euphorbia nicaeensis* All. leaves and stems were performed using standardized laboratory protocols. The presence or absence of phenols, tannins ^[15], flavonoids ^[16], saponins ^[17], flavonol, carbohydrates, alkaloids, gums, amino acids ^[18], sterols and polyterpenes ^[19], was analyzed accordingly.

2.4. Total phenolic content (TPC)

The phenolic compounds were measured using Folin-Ciocalteu method [20]. In brief, 50 μ l of Folin-Ciocalteu reagent was added to the sample, followed by 150 μ l of Na_2CO_3 after 10 min, and the volume was made up to 1 ml with water. The absorbance was measured in a spectrophotometer reader at 760 nm after 2 hours of incubation and compared to the Gallic acid calibration curve (5 – 100 μ g/ml), $R^2= 0.998$. The results were expressed as mg Gallic acid equivalents /g dry weight (mg GAE/g DW).

2.5. Total flavonoid content (TFC)

According to [21], 250 μ l of the extract solution was mixed with 1 ml of distilled water and 75 μ l of NaNO_2 solution (5 %). After 6 min, 75 μ l of AlCl_3 solution (10 %) was added. The mixture was allowed to stand for 6 min before adding 1 ml of NaOH solution (4 %) and bringing the final volume to 2.5 ml with distilled water. The mixture was properly vortexed and allowed to stand for 15 min in darkness. The absorbance was measured at 510 nm and compared to the catechin standard curve (5 – 400 μ g/ml) $R^2= 0.997$. The results were expressed as mg of catechin equivalent per g dry weight (mg CE/g DW).

2.6. Flavonol content (FC)

The content of flavonols was determined according to the method described by [22]. An aliquot of 500 μ l of extract was added to 500 μ l of Aluminium chloride (20 mg/ml) and 1.5 ml of sodium acetate (50 mg/ml). The mixture was properly vortexed and left to stand in darkness for 2.5 h. The absorbance was read at 440 nm, and the result was expressed referring to the absorbance of standard quercetin solution prepared in the same conditions (5 – 200 μ g/ml), $R^2= 0,995$. The flavonol content is expressed in mg Quercetin equivalents per g dry weight (mg QE/g DW).

2.7. Condensed tannin content (CTC)

The assay was executed as reported previously [23]. An aliquot of 100 μ l of each extract was added to 1.5 ml of Vanillin (4 %) and 750 μ l of concentrated hydrochloric acid (HCL). The mixture was vortexed, left to stand in the dark for 20 min, and the absorbance was recorded at 500 nm. The condensed tannin content was counted related to the Catechin calibration curve (5 – 400 μ g/ml) $R^2= 0.985$, elaborated in the same manner. The results are expressed as mg of Catechin equivalent per 100 g dry weight (mg CE/100g DW).

2.8. Lipid-soluble pigment content

According to [21], 150 mg of vegetal powder (leaves and stems) was mixed for 1 min with 10 ml of Acetone/Hexane (4:6). The mixture was filtered through Whatman filter paper grade 4. The absorbance was recorded at 453,505,645, and 663 nm. The content of β -carotene, lycopene, chlorophyll a, and chlorophyll b was calculated according to the following equations, expressed in μ g per g dry weight (DW):

$$\beta\text{-carotene} = 0.216 \times A663 - 1.220 \times A645 - 0.304 \times A505 + 0.452 \times A453$$

$$\text{Lycopene} = -0.0458 \times A663 + 0.204 \times A645 - 0.304 \times A505 + 0.452 \times A453$$

$$\text{Chlorophyll a} = 0.999 \times A663 - 0.0989 \times A645$$

$$\text{Chlorophyll b} = 0.328 \times A663 + 1.77 \times A645$$

A453, A505, A645, and A663 are the absorbance measured at 453, 505, 645, and 663 nm, respectively.

2.9. *In vitro* antioxidant activity

2.9.1. 2,2-Diphenyl-1-picrylhydrazyl (DPPH) radical-scavenging assay

The radical scavenging ability of the extract was carried out as described by [24–26]. 500 μ l of extract solution was mixed with 1 ml of DPPH solution (0.1 mM in methanol) freshly prepared. The mixture was vortexed properly and left to stand in the dark for 60 min. The reduction of DPPH was measured at 515 nm, and the scavenging effect was estimated based on the percentage of scavenging DPPH radicals using the following equation:

$$\% \text{ DPPH scavenging activity} = \frac{ADPPH - A_s}{ADPPH} \times 100$$

ADPPH is the absorbance of DPPH and AS is the absorbance of DPPH when the sample has been added at different concentrations. The IC₅₀ value is the concentration that scavenges 50 % of DPPH radicals, and it is calculated from the scavenging effect percentage graph. Ascorbic acid was used as a standard.

2.9.2. Scavenging of hydrogen peroxide assay

The performance of hydrogen peroxide (H₂O₂) scavenging was investigated as follows [27]. The phosphate buffer (50 mM, 7.4 pH) received a 40 mM H₂O₂ solution. In order to quantify the mixture's absorbance spectrophotometrically at 230 nm, all experimental samples were combined with 0.6 ml of H₂O₂ solution. The mixture was then incubated for 10 minutes. Ascorbic acid served as the standard, while phosphate buffer functioned as the control. Hydrogen peroxide scavenging (%) was calculated using the formula below.

$$\% \text{ Hydrogen peroxide scavenging activity} = \frac{A_0 - A_s}{A_0} \times 100$$

Where A_s is the absorbance in the presence of the ascorbic acid standard or samples, and A₀ is the absorbance of the blank. Ascorbic acid was used as a standard.

2.10. *In vitro* anti-inflammatory activity

2.10.1. Bovine Serum Albumin Assay (BSA)

Protein denaturation is one of the causes of inflammatory and rheumatic diseases, as reported by a previous study [28]. Any compound that provides greater than 20% inhibition of protein denaturation is considered a potential anti-inflammatory agent and could be useful for treating several diseases [29].

BSA assay of leaf and stem extracts was determined using the method reported by [27]. 0.45 ml of bovine serum albumin was mixed with 0.05 ml of samples at various concentrations (200 - 2000 µg/ml). The mixture was incubated for 25 min at 40°C, and phosphate buffer saline (2.5 ml; pH 6.3) was added to tubes. The control received phosphate buffer solution (0.05 ml) instead of extract. The absorbance was measured using a spectrophotometer at 660 nm, and the percentage inhibition of BSA denaturation was calculated using the following equation:

$$\% \text{ inhibition of BSA denaturation} = \frac{A1 - A2}{A1} \times 100$$

Where A1 = absorbance of the control and A2 = absorbance of the test sample.

2.10.2. Chicken Egg albumin assay (CEA)

Using the technique described by [30], the ability of our extracts to inhibit protein denaturation was examined. To 2.8 ml of PBS solution, 2 ml of extract and 0.2 ml of chicken egg albumin were added. All samples were kept at room temperature for 15 minutes before being heated to 70 °C for 10 minutes. Diclofenac was employed as standard, and the absorbance was measured at 660 nm. The equation below was used to determine the inhibition of egg albumin denaturation:

$$\% \text{ inhibition of egg albumin denaturation} = \frac{A1 - A2}{A1} \times 100$$

A1 is the control's absorbance and A2 is the test sample's absorbance.

2.11. Phenolic profile analysis by HPLC-UV-MS/MS

Phenolic compounds were qualitatively analyzed using a liquid chromatography system coupled with a triple quadrupole mass spectrometer (Thermo Fisher Scientific, San Jose, CA, USA). The suggested technique was carried out using a Kinetex C18 reversed-phase column (100 x 4.6 mm, 2.6 µm particles). From solvent A (0.1% formic acid aqueous solution) and solvent B (methanol), gradient separation was created following the method described by [31]. Retention time and spectrum matching with nine standards were used to identify the phenolic compounds, along with the NIST-MS/MS library.

2.12. Statistical analysis

T-test was used to evaluate the statistical differences between extraction methods (maceration and probe-assisted ultrasound extraction) and organs (leaves and stems). Experiments were run in triplicate, and the results are expressed as mean values of three analyses. Data are statistically significant at $p < 0.05$ and highly significant at $p < 0.001$. Pearson correlation coefficient (r), DPPH (IC_{50}), and H_2O_2 (IC_{50}) were determined using GraphPad prism 8.0.2.

3. Results

3.1. Phytochemical screening

The results of phytochemical screening of *Euphorbia nicaeensis* All. leaves and stems are shown in Table 1. Phenols, tannins, saponins, and carbohydrates are strongly present in all plant organs. Flavonoids, flavonol glycosides, sterols, and polyterpenes are highly present in leaves and small amounts in stems. Alkaloids and amino acids are present in tiny quantities, quinones and mucilage are absent in all plant organs.

Table 1. Phytochemical screening results of leaves and stems of *Euphorbia nicaeensis* All.

Phytochemical compounds	Leaves	Stems
Phenols / Tanins	(++)	(++)
Flavonoids	(++)	(+)
Flavonol glycosides	(+)	(+/-)
Alkaloids	(+/-)	(+/-)
Saponins	(++)	(++)
Carbohydrates	(++)	(++)
Quinones	(-)	(-)
Gums and mucilages	(-)	(-)
Amino acids	(+)	(+)
Sterols and Polyterpenes	(++)	(+)

(+): Presence of phytochemicals, (-): Absence of phytochemicals, (++): Presence in significant amounts, (+/-): Presence in small amounts.

3.2. Total phenol (TPC), Total flavonoid (TFC), Flavonol content (FC), and Condensed tannin content (CTC)

The carried-out tests indicated that the leaves and stems of *Euphorbia nicaeensis* are rich in phenolic compounds. The percentage yield of leaf and stem extracts by maceration (3.32%, 1.68%), and ultrasonic-assisted extraction ($12.69 \pm 0.17\%$, $4.62 \pm 1.55\%$), respectively, showed that the highest extraction yield of both leaves and stems was obtained with probe ultrasonic-assisted extraction (PUAE). The (TPC) values of PUAE extract of leaves and stems (80.13 ± 1.61 mg GAE/g DW, 77.79 ± 2.29 mg GAE/g DW), respectively, were statically higher (t-test, $p < 0.05$) than that of maceration extracts (68.69 ± 0.91 mg GAE/g DW, 68.31 ± 1.83 mg GAE/g DW). This data also reveals that the leaves are richer in flavonoids, condensed tannins, and flavonols than the stems as presented in Table 2.

Table 2. Total phenol, flavonoids, condensed tannins, flavonols contents and extraction yield of maceration extracts and probe ultrasonic-assisted extract of *Euphorbia nicaeensis* All. leaves and stems.

	ELM*	ESM*	ELU*	ESU*	p_1	p_2	p_3	p_4
TPC (mg GAE/g DW)	68.69 ± 0.91	68.31 ± 1.83	80.13 ± 1.61	77.79 ± 2.29	0.574	0.042	<0.001	<0.001
TFC (mg CE/g DW)	32.83 ± 3.84	21.83 ± 9.93	49.22 ± 0.192	45.27 ± 2.5	0.047	0.031	0.004	0.010
CTC (mg CE/100gDW)	1.33 ± 1.1	1.88 ± 1.92	7.61 ± 1.92	1.22 ± 0.92	0.450	0.002	0.001	0.361
FC (mg QE/g DW)	4.80 ± 0.08	1.23 ± 0.27	36.14 ± 0.08	5.88 ± 0.33	<0.001	<0.001	<0.001	<0.001
Extraction yield (%)	3.32 ± 0.59	1.68 ± 0.28	12.69 ± 0.17	4.62 ± 1.55	0.024	0.011	<0.001	0.077

ELM: Euphorbia leaf maceration, ESM: Euphorbia stem maceration, ELU: Euphorbia leaves ultrasonic-assisted extraction, ESU: Euphorbia stems ultrasonic-assisted extraction, TPC: Total flavonoid content, TFC: Total flavonoid content, CTC: Condensed tannin contents, FC: Flavonols content, p_1 : p -value (ELM – ESM), p_2 : p -value (ELU – ESU), p_3 : p -value (ELM – ELU), p_4 : p -value (ESM – ESU). *Means \pm SD from triplicate determinations. p -value is considered significant at $p < 0.05$ and highly significant at $p < 0.001$.

3.3. Lipid-soluble pigment content

As shown in Figure 2, this study revealed that leaves and stems differed significantly ($p < 0.05$) in the content of chlorophyll pigment. The leaves are richer in chlorophyll a and b ($0.167 \pm 0.001 \mu\text{g/g DW}$, $0.108 \pm 0.002 \mu\text{g/g DW}$) than the stems ($0.016 \mu\text{g/g DW}$, $0.049 \pm 0.001 \mu\text{g/g DW}$). These findings also showed that leaves have significantly higher ($p < 0.05$) β -carotene content ($0.035 \pm 0.001 \mu\text{g/g DW}$) compared to stems ($0.006 \pm 0.001 \mu\text{g/g DW}$). In addition, low lycopene concentration was detected in both leaves and stems ($0.033 \pm 0.001 \mu\text{g/g DW}$, $0.036 \mu\text{g/g DW}$), respectively, with no significant difference ($p > 0.05$).




Figure 2. Lipid-soluble pigment content of leaves and stems of *Euphorbia nicaeensis*. Values are means ± S.D of three independent measurements. EL: Euphorbia leaves, ES: Euphorbia stems. The values with the same superscript letters are not significantly different ($p > 0.05$).

3.4. In vitro antioxidant activity

The evaluation of the antioxidant activity of *E. nicaeensis* leaf and stem extracts was screened by 2,2-Diphenyl-1-picrylhydrazyl (DPPH) assay and scavenging of hydrogen peroxide (H_2O_2) assay. All extracts showed antioxidant activity in Figure 3. The results were positively correlated to the concentration, and the maximum scavenging activity was recorded at the highest concentration for all extracts. Furthermore, it was observed that PUAE provided a strong significant DPPH scavenging effect ($p < 0.01$), translated by a low IC_{50} value of leaf extracts ($\text{IC}_{50} = 23.47 \pm 0.62 \mu\text{g/ml}$) closely compared to Ascorbic Acid (IC_{50}

=19.89±0.14 µg/ml). While the maceration leaf extract showed a low antioxidant effect (IC₅₀ = 79.76±1.93 µg/ml).




Figure 3. DPPH radical scavenging activity (A) and H₂O₂ scavenging activity (B) versus concentration (µg/mL). ELM: Euphorbia leaf maceration, ESM: Euphorbia stem maceration, ELU: Euphorbia leaf ultrasonic-assisted extraction, ESU: Euphorbia stem ultrasonic-assisted extraction. The results are expressed as means ± SD of three independent measurements.

The stem extract of PUAE also exhibited a moderate antioxidant effect (IC₅₀ = 42.45 ± 2.30µg/ml) compared to the maceration extract (IC₅₀ = 61.16 ± 1.53µg/ml). However, it was observed that stem maceration extract has a significantly (*p*<0.01) lower IC₅₀ value than leaf maceration extract, which implies a higher antioxidant capacity, as shown in Table 3, similarly, for H₂O₂ scavenging results. The lowest IC₅₀ was observed in PUAE leaf extract (IC₅₀=110.27 ± 3.59 µg/ml) followed by stem extract (IC₅₀ = 131.51 ± 1.80 µg/ml) in comparison to maceration (IC₅₀ = 678.32 ± 7.22 µg/ml, IC₅₀ = 1073.63 ± 1.08 µg/ml, respectively).

Table 3. Scavenging activity of leaf and stem extracts of *Euphorbia nicaeensis* All.

	Scavenging activity (IC ₅₀ µg/ml)	
	DPPH assay	H ₂ O ₂ assay
ESM*	61.16 ± 1.53	1073.63 ± 1.08
ELM*	79.76 ± 1.93	678.32 ± 7.22
ESU*	42.46 ± 2.30	131.51 ± 1.80
ELU*	23.48 ± 0.62	110.27 ± 3.59
AA*	19.90 ± 0.14	35.41 ± 0.23

*Means ± SD from triplicate determinations, *p* < 0.001.

The correlation analysis between antioxidant activity and phenolic content of the extracts is presented in Table 4. Pearson's coefficient revealed a positive relationship between antioxidant activity and total phenolics and flavonoids' plant content. The DPPH assay of antioxidant activity and TPC showed a Pearson coefficient of $r=0.8827$ and $r=0.7768$ for total flavonoids content with no significant correlation ($p>0.05$). Moreover, a highly significant correlation was observed between the H_2O_2 assay and the composition of the plant in both TPC ($p < 0.001$) and TFC ($p < 0.05$).

Table 4. Pearson correlation test between total phenolic content, total flavonoids content, and antioxidant activity.

Pearson's correlation (r)		
	DPPH (1/IC ₅₀)	H ₂ O ₂ (1/IC ₅₀)
TPC	0.8827	0.9995**
TFC	0.7768	0.9505*

*Significant at $p < 0.05$; **Significant at $p < 0.001$; DPPH: DPPH scavenging activity; H₂O₂: hydrogen peroxide scavenging activity; TPC: Total flavonoid content; TFC: Total flavonoid content.

3.5. *In vitro* anti-inflammatory activity

According to the results presented in Table 5, all extracts showed an inhibition of Bovine Serum Albumin (BSA) and chicken egg albumin denaturation (CEA), which increased with the sample concentration. The highest inhibitions were observed at the 2000 µg/ml dose for all extracts.

The findings revealed a notable inhibition of serum albumin denaturation (83.98% and 75.39%, $p < 0.001$) and egg albumin denaturation (82.04% and 78.87%, $p < 0.001$) induced by PUAE extracts of stems and leaves, respectively, at the dose of (2000 µg/ml). The maceration extracts of leaves and stems showed a lower inhibition of bovine serum albumin (38.28% and 43.36%, $p < 0.001$) at the same concentration. The inhibition of egg albumin revealed similar results. Maceration extracts from stems and leaves showed a weak inhibition (49.85%, 62.52%, respectively). While the standard drug diclofenac sodium exhibited potent inhibition (97.73%, $p < 0.001$) at the same concentration when compared to the control.

Stem extracts from maceration and ultrasound-assisted extraction showed a significantly higher level of protein denaturation inhibition ($p < 0.05$) than leaf extracts at the concentration range of (2000- 500 $\mu\text{g/ml}$).

Table 5. In vitro anti-inflammatory activity of *Euphorbia nicaeensis* All.

Treatment	Dosage ($\mu\text{g.ml}^{-1}$)	Bovine serum albumin denaturation		Chicken Egg albumin denaturation	
		Absorbance	% Inhibition	Absorbance	% Inhibition
Control	-	0.917 ± 0.015	-	0.913 ± 0.015	-
ELM	200	0.652 ± 0.0006	28.90	0.286 ± 0.003	72.48
	300	0.591 ± 0.002^a	35.54	0.221 ± 0.001	72.08
	500	0.571 ± 0.0026	37.77	0.218 ± 0.003	70.95
	1000	0.570 ± 0.0021	37.89	0.197 ± 0.001	68.80
	2000	0.566 ± 0.001	38.28	0.193 ± 0.001	62.52
ESM	200	0.677 ± 0.0072	26.17	0.279 ± 0.001	73.32
	300	0.570 ± 0.006	37.89	0.223 ± 0.001	63.94
	500	0.541 ± 0.0066	41.02	0.206 ± 0.004	59.82
	1000	0.536 ± 0.0061	42.19	0.192 ± 0.001	52.04
	2000	0.519 ± 0.009	43.36	0.164 ± 0.004	49.85
ELU	200	0.638 ± 0.003	30.47	0.342 ± 0.003	68.72
	300	0.605 ± 0.0096^a	33.98	0.285 ± 0.004^b	75.77
	500	0.390 ± 0.0046	57.42	0.265 ± 0.003	76.13
	1000	0.276 ± 0.0053	69.92	0.255 ± 0.004	78.43
	2000	0.226 ± 0.001	75.39	0.251 ± 0.001	78.87
ESU	200	0.613 ± 0.0044	33.20	0.458 ± 0.006	69.49
	300	0.537 ± 0.001	41.41	0.438 ± 0.012^b	75.62
	500	0.308 ± 0.0053	66.41	0.367 ± 0.002	77.48
	1000	0.208 ± 0.000	77.34	0.329 ± 0.003	78.98
	2000	0.147 ± 0.0035	83.98	0.244 ± 0.002	82.04
Diclofenac sodium	200	0.079 ± 0.0026	91.41	0.054 ± 0.004	94.12
	300	0.062 ± 0.001	93.23	0.047 ± 0.002	94.89
	500	0.057 ± 0.0036	93.75	0.041 ± 0.002	95.47
	1000	0.050 ± 0.0017	94.53	0.035 ± 0.004	96.17
	2000	0.021 ± 0.0044	97.73	0.022 ± 0.004	97.59

The results are expressed as means \pm SD of three independent measurements. ELM: Euphorbia leaf maceration, ESM: Euphorbia stem maceration. ELU: *Euphorbia* leaf ultrasonic-assisted extraction, ESU: *Euphorbia* stem ultrasonic-assisted extraction. the values with the same superscript letters are not significantly different ($p > 0.05$).

3.6. Phenolic profile analysis by HPLC-MS




Figure 4. HPLC-MS/MS chromatogram of UAE stem extract.




Figure 5. HPLC-MS/MS chromatogram of UAE leaf extract.

Based on the previous results of the phenolic compounds assay, UAE leaf, and stem extracts were retained for HPLC-UV-MS/MS analysis. The presence of the compounds identified by HPLC-UV analysis was confirmed using HPLC-MS/MS. Comparisons with reference standards, NIST library, and earlier literature reports helped identify. The retention times of the standards tested did not match the peaks detected in the chromatograms of the leaf and stem extracts (Figures 4 and 5). This is because many are in the form of sugar derivatives that are rarely available. Therefore, mass spectrometry was utilized to characterize the peaks and allowed further identification of the compounds. Comparing the MS/MS data with those in the NIST library and other references revealed several phenolic compounds.

Table 6 demonstrates the abundance of different phytochemicals present in both extracts, with a slight difference in composition, including flavonols, phenolic acids, lignan, calchone, carotenoids, and alkaloids. Fargesin, 2'-hydroxy-3,4,4',6'- tetramethoxychalcone, nuciferin, quercetin 3'-methyl ether, and isoquercetin are detected in both extracts with different peak airs. Isoquercetin was detected as the predominant compound in the leaf extract, followed by quercitrin hydrate and guaiaverin. The stem extract is rich in 2'-hydroxy-3,4,4',6'- tetramethoxychalcone, representing the highest peak (22.04%), followed by Quercetin 3'-methyl ether.

Table 6. HPLC-MS/MS tentative identification of phenolics and derivatives in *E. nicaeensis* leaf and stem extracts.

Leave					Stem				
N°	RT	Tentative identification	Air %	Ref	N°	RT	Tentative identification	Air%	Ref
1	1.70	Fargesin	5.72	N	1	1.70	Fargesin	3.34	N
2	1.92	2'-hydroxy-3,4,4',6'-tetramethoxychalcone	4.51	N	2	1.96	2'-hydroxy-3,4,4',6'-tetramethoxychalcone	22.04	N
3	3.18	Nuciferine	2.08	N	3	3.18	<u>Nuciferine</u>	8.14	<u>N</u>
4	4.86	Quercetin 3'-methyl ether	3.90	N	4	4.78	Quercetin 3'-methyl ether	10.02	N
5	10.08	5,7,3',4'-Tetramethoxyisoflavone	2.11	N	5	10.38	Lutein	3.65	N
6	10.97	Galloylquinic acid	1.79	[32]	6	17.77	Dilinolenin	3.11	N
7	19.90	Quercetin-galactoside-gallate	8.70	[33]	7	21.11	Isoquercetine	0.94	N
8	20.47	Myricitrin	6.59	N	8	21.52	2'-Hydroxy-2,4,4'-trimethoxychalcone	9.59	N
9	21.11	Isoquercetin	27.49	N					
10	22.57	Guaiaverin	11.78	N					
11	22.99	Quercitrin hydrate	14.49	[31]					

RT: retention time, N: NIST-MS/MS library, Air %: Relative peak area

4. Discussion

Most of the phytochemical screening findings align with a previous investigation conducted in Serbia for *Euphorbia nicaeensis* ssp. *glareosa* [34]. However, some differences were observed, namely, the presence of alkaloids and the absence of quinones in *E. nicaeensis* All. leaves and stems, compared to this subspecies. This could be due to several parameters, including genetics, differences in harvesting site, rainfall, light, season, harvesting period, topography, soil type, and extraction method [35].

The phytochemical assay showed that the leaves and stems of *E. nicaeensis* are rich in total phenolic, flavonoids, and flavonols and contain a moderate amount of condensed tannins. Flavonoids, particularly flavonols, are effective antifungal agents against various pathogens, including *Candida albicans*, *Candida glabrata*, *Candida krusei*, *Candida parapsilosis*, *Candida tropicalis*, *Trichophyton rubrum* and *Trichophyton beigeli*. [36–40]. Condensed tannins have also been demonstrated to have potent anthelmintic properties against worm infections. These findings support the traditional usage of *Euphorbia nicaeensis* to treat fungal infections and parasite disorders [41]. By comparing the results of total phenolic content (TPC), total flavonoid content (TFC), condensed tannin content (CTC), flavonol content (FC), and the yield of the two extraction methods, it can be concluded that probe ultrasound-assisted extraction (PUAE) is more efficient than maceration. These might be linked to acoustic cavitation processes, which could cause a forceful impact on the solid surface, resulting in an enhanced extraction rate, as already reported [42]. It is therefore inferred that PUAE could be a convenient technique for phenolic compound extraction. However, the maceration extract of stems proved to have a higher antioxidant effect than leaf maceration extract. These results can be interpreted by the variation of the nature and chemical structure of the phenolic compounds present in each sample. In some cases, the highest antioxidant activity may be observed in extracts with a low phenolic content. Synergy may occur between the major antioxidants (phenolic compounds) and other minor constituents of the plant, so this could significantly impact the differences in their antioxidant activity [43].

This study revealed that *E. nicaeensis* All. leaves possess a potent antioxidant effect close to ascorbic acid activity. A higher free radical scavenging and hydrogen peroxide scavenging effect were observed in probe ultrasound-assisted extraction leaf extracts, compared to stems. The difference between leaves and stems in extraction yield and total phenolic contents could partially explain this. Studies have reported that leaves contain a high amount of phenolic compounds than stems [44]. A potent correlation was observed between phenolic compounds and H₂O₂ scavenging activity, as reported by reports [45,46]. However, the inclusion of other substances, particularly sugars, which, according to [47], can interfere with quantification and antioxidant activity assays, may cause a non-significant correlation between TPC and DPPH.

HPLC-UV-MS/MS analysis showed that the tested extracts are rich in phenolic compounds, especially flavonoids (flavonol glycosides). Quercetin and its derivatives are the

most abandoned compounds in the tested extracts, especially in the leaves, and studies have reported their different biological virtues ^[48,49]. The presence of isoquercetin and quercitrin hydrate represented by the highest peaks, may be among the leading causes of the significant antioxidant properties of the leaves. Fargesin, myricitrin, and guaiaverin also play a crucial role in plant health and display several therapeutic values due to their antioxidant or/and anti-inflammatory properties ^[50,51].

From this, we can infer that the phenolic compounds of the plant are the major cause of the scavenging effect of the extracts, supporting the results obtained and explaining why the leaves have a higher antioxidant power than the stems. Therefore, leaves are a potential source of natural antioxidants and bioactive compounds.

The results of previous research on different species and subspecies of the Euphorbiaceae family have demonstrated that *E. nicaeensis* All. has a higher DPPH radical inhibition potential than *E. retusa* Forssk. (IC₅₀ leaf = 287.52±2.92 µg/ml, IC₅₀ stem=225.87±3.88 µg/ml ^[52] and *E. hirta* L. (IC₅₀ leaf = 803 µg/ml, IC₅₀ stem= 1358 µg/ml) ^[53], as well as *E. heterophylla* (IC₅₀ = 194.28±0.22 µg/ml) ^[54]. The subspecies *Euphorbia nicaeensis* ssp. *glareosa* showed more than 50% inhibition of DPPH (56.5%), which qualifies the plant as moderately active ^[34]. Yet no previous study on the in-vitro antioxidant activity of *E. nicaeensis* All. has been published.

The aerial part of *E. nicaeensis* All. is recognized as having anti-inflammatory virtue as it contains glyceroglycolipids tested for their anti-inflammatory activity ^[55]. This study's results agree with what has been previously published. In vitro tests showed that the plant has a strong anti-inflammatory effect, inhibiting the degradation of bovine serum albumin (BSA) and chicken egg albumin (CEA), observed mainly in the stem extracts. Nuciferine (alkaloid) and fargesin (lignan) are among the main compounds identified in the stem extract according to chromatograms and MS/MS identification; they are phytochemical compounds known for their anti-inflammatory virtues ^[50,56] which proves the results of the anti-inflammatory activities performed.

The inhibition of thermal degradation of proteins by extract can be explained by several mechanisms, depending on the type of plant extract and the protein studied ^[57]: Proteins can be protected from thermal degradation by antioxidants because they can scavenge reactive oxygen species (ROS) and reduce oxidative degradation ^[58]. When phenolic chemicals interact with proteins, their properties, such as solubility, digestibility, and thermal stability, may alter. However, extracts rich in phenolic compounds have been employed to inhibit protein denaturation and improve thermal stability ^[59]. The antioxidant effect of *E. nicaeensis* extracts resulting from the presence of phenolic compounds, specifically quercetin, and derivatives, play an essential role in the stability of proteins and inhibit denaturation caused by heat treatment, as already reported.

Based on the results of anti-inflammatory activity and those reported by a previous study, we can deduce that the aerial part of *E. nicaeensis*, especially the stems, provides a

potent anti-inflammatory activity and could be considered a potential anti-inflammatory agent.

5. Conclusion

This study aimed to investigate the phytochemical profile and determine the antioxidant and anti-inflammatory activities of leaf and stem extracts of *Euphorbia nicaeensis* All. It has been observed that ultrasound-assisted extraction is an efficient and recommended method for phenolic compound extraction. This work also showed that all extracts are rich in phenolic compounds and exhibited efficient antioxidant and anti-inflammatory properties with significant differences between organs.

HPLC-UV-MS/MS confirmed the presence of phenolic compounds, particularly quercetin and its derivatives, present mainly in the leaf extracts. The leaves showed the highest antioxidant activity, while the most potent anti-inflammatory effect was observed in the stem extracts. The phenolic composition indicated the presence of several compounds with anti-inflammatory properties. These findings confirmed *E. nicaeensis* historical medicinal uses and opened up new ways and possibilities of developing this plant in the pharmaceutical and food fields as a new source of bioactive compounds unaffected by commercial breeding. However, it is evident that additional research, such as the molecular identification of the bioactive substances responsible for these biological activities, is required to offer a more satisfactory and clearer perspective to this study.

Author Contributions: Conceptualization, KB and HT.; methodology, KB, HT, CE, RS; software, NEH, CE.; validation, X.X., Y.Y. and Z.Z.; formal analysis, NEH, HNM, KB, KM, MH, RF; investigation KB, HT, CE; resources KWG, AB, HNM; data curation, AK and KB.; writing—original draft preparation, KB and CE; writing—review and editing AB, NEH, HT, KWG.

Funding: No external funding was provided for this research.

Acknowledgments: Authors of this work express their sincere thanks to everyone who contributed to the realization of this study, especially Abderrahmane Belhouari, Professor of statistics at the Faculty of Science BEN MSIK and Said Zaidoune, Professor of English at the Faculty of Letters BEN MSIK.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Garg AK, Faheem M, and Singh S. Role of Medicinal Plant in Human Health Disease. *Asian J Plant Sci* 2021; 11(1). doi:10.36648/2249-7412.11.1.19-21
2. Elachouri M, Kharchoufa L, Fakchich J, *et al.* Ancestral phytotherapeutic practices in Morocco: regards on history, current state, regulatory and safety of commonly used herbal medicine. *Arab J Chem* 2021; 8(1): 133-149.
3. Samman NA, Martin A, and Puech S. Inflorescence architecture variability and its possible relationship to environment or age in a Mediterranean species, *Euphorbia nicaeensis* All. (*Euphorbiaceae*). *Bot J Linn* 2001; 136(1): 99-105. doi:10.1111/j.1095-8339.2001.tb00558.x
4. Bettar I, González-Miret ML, Hernanz D, *et al.* Characterisation of Moroccan Spurge (*Euphorbia*) honeys by their physicochemical characteristics, mineral contents and colour. *Arab J Chem* 2019; 12(8): 2052-2060.
5. Bellakhdar J. Contribution à l'étude de la pharmacopée traditionnelle au Maroc: la situation actuelle, les produits, les sources du savoir (enquête ethnopharmacologique de terrain réalisée de 1969 à 1992).; 1997.

6. Fakchich J and Elachouri M. An overview on ethnobotanico-pharmacological studies carried out in Morocco, from 1991 to 2015: Systematic review (part 1). *J Ethnopharmacol* 2021; 267: 113200. doi:10.1016/j.jep.2020.113200
7. Chow YL and Tham HW. Methicillin-resistant *Staphylococcus aureus* (MRSA) on dispensing counters of community pharmacies in Klang Valley. *Prog Microbes Mol Biol* 2020; 3(1).
8. Lau AWY, Tan LTH, Ab Mutalib NS, *et al.* The chemistry of gut microbiome in health and diseases. *Prog Microbes Mol Biol* 2021; 4(1).
9. Krstić G, Jadranin M, Todorović NM, *et al.* Jatrophone diterpenoids with multidrug-resistance modulating activity from the latex of *Euphorbia nicaeensis*. *Phytochemistry* 2018; 148: 104-112. doi:10.1016/j.phytochem.2018.01.016
10. Krstić GB, Kostić A, Jadranin MB, *et al.* Two new jatrophone diterpenes from the roots of *Euphorbia nicaeensis*. *J Serb Chem* 2021; 86(12): 1219-1228.
11. Özbigin S and Çitoğlu G. Uses of some *Euphorbia* species in traditional medicine in Turkey and their biological activities. *Turkish J Pharm* 2012; 9: 241-256.
12. Cateni F, Falsone G, Zilic J, *et al.* Glyceroglycolipids from *Euphorbia nicaeensis* All. with antiinflammatory activity. *Zhdankin VV, ed. Arkivoc* 2004; 2004(5): 54-65. doi:10.3998/ark.5550190.0005.506
13. Shi QW, Su XH, and Kiyota H. Chemical and Pharmacological Research of the Plants in Genus *Euphorbia*. *Chem Rev* 2008; 108(10): 4295-4327. doi:10.1021/cr078350s
14. Ranjha MMAN, Irfan S, Lorenzo JM, *et al.* Sonication, a Potential Technique for Extraction of Phytoconstituents: A Systematic Review. *Processes* 2021; 9(8): 1406. doi:10.3390/pr9081406
15. Yadav RNS and Agarwala M. Phytochemical analysis of some medicinal plants. *J Phytol* Published online December 14, 2011. Accessed October 22, 2022. <https://updatepublishing.com/journal/index.php/jp/article/view/2737>
16. Pandey A and Tripathi S. Concept of standardization, extraction and pre phytochemical screening strategies for herbal drug. *J Pharmacogn Phytochem* 2014; 2(5): 115-119.
17. Yadav M, Chatterji S, Gupta SK, *et al.* Preliminary phytochemical screening of six medicinal plants used in traditional medicine. *Int J Pharm* 2014; 6: 539-542.
18. Parimelazhagan T. *Pharmacological Assays of Plant-Based Natural Products*. Vol 71. Springer International Publishing; 2016. doi:10.1007/978-3-319-26811-8
19. Koffi J, Bla B, Yapi HF, *et al.* Phytochemical screening of some medicinal plants in Côte D'ivoire and evaluation of their extraction efficiency. *Int J Pharm* 2015; 7: 563-569.
20. Elouafy Y, Mortada S, El Yadini A, *et al.* Bioactivity of Walnut: Investigating the Triterpenoid Saponin Extracts of *Juglans regia* Kernels for Antioxidant, Anti-diabetic, and Antimicrobial Properties. *Prog Microbes Mol Biol* 2023; 6(1).
21. Barros L, Cabrita L, Boas MV, *et al.* Chemical, biochemical and electrochemical assays to evaluate phytochemicals and antioxidant activity of wild plants. *Food Chem* 2011; 127(4): 1600-1608. doi:10.1016/j.foodchem.2011.02.024
22. Awah FM, Uzoegwu PN, Ifeonu P, *et al.* Free radical scavenging activity, phenolic contents and cytotoxicity of selected Nigerian medicinal plants. *Food Chem* 2012; 131(4): 1279-1286. doi:10.1016/j.foodchem.2011.09.118
23. Salar RK, and Purewal SS. Improvement of DNA damage protection and antioxidant activity of biotransformed pearl millet (*Pennisetum glaucum*) cultivar PUSA-415 using *Aspergillus oryzae* MTCC 3107. *Biocatal Agric Biotechnol* 2016; 8: 221-227. doi:10.1016/j.bcab.2016.10.005
24. Kemung HM, Tan LTH, Chan KG, *et al.* *Streptomyces* sp. strain MUSC 5 from mangrove forest in Malaysia: Identification, antioxidant potential and chemical profiling of its methanolic extract. *Prog Microbes Mol Biol* 2020; 3(1).

25. El Hachlafi N, Chebat A, Fikri-Benbrahim K. Ethnopharmacology, Phytochemistry, and Pharmacological Properties of *Thymus satureioides* Coss. *Evid Based Complement Altern Med* 2021; 2021.
26. Benkhaira N, Koraichi SI, and Fikri-Benbrahim K. In vitro methods to study antioxidant and some biological activities of essential oils: a review. *Biointerface Res Appl Chem* 2022; 12(3): 3332.
27. Qamar M, Akhtar S, Ismail T, *et al.* *Syzygium cumini*(L.), Skeels fruit extracts: In vitro and in vivo anti-inflammatory properties. *J Ethnopharmacol* 2021; 271: 113805. doi:10.1016/j.jep.2021.113805
28. Kar B, Kumar RS, Karmakar I, *et al.* Antioxidant and in vitro anti-inflammatory activities of *Mimusops elengi* leaves. *Asian Pac J Trop* 2012; 2(2, Supplement): S976-S980. doi:10.1016/S2221-1691(12)60346-3
29. Gondkar AS, Deshmukh VK, and Chaudhari SR. Synthesis, characterization and in-vitro anti-inflammatory activity of some substituted 1,2,3,4 tetrahydropyrimidine derivatives. *Drug Invent Today* 2013; 5(3): 175-181. doi:10.1016/j.dit.2013.04.004
30. Kola P, Metowogo K, Manjula SN, *et al.* Ethnopharmacological evaluation of antioxidant, anti-angiogenic, and anti-inflammatory activity of some traditional medicinal plants used for treatment of cancer in Togo/Africa. *J Ethnopharmacol* 2022; 283: 114673. doi:10.1016/j.jep.2021.114673
31. Puigventós L, Navarro M, Alechaga É, *et al.* Determination of polyphenolic profiles by liquid chromatography-electrospray-tandem mass spectrometry for the authentication of fruit extracts. *Anal Bioanal Chem* 2015; 407(2): 597-608. doi:10.1007/s00216-014-8298-2
32. Clifford MN, Stoupi S, Kuhnert N. Profiling and characterization by LC-MSn of the galloylquinic acids of green tea, tara tannin, and tannic acid. *J Agric Food Chem* 2007; 55(8): 2797-2807. doi:10.1021/jf0635331
33. Pereira P, Cebola MJ, Oliveira MC, *et al.* Antioxidant capacity and identification of bioactive compounds of *Myrtus communis* L. extract obtained by ultrasound-assisted extraction. *J Food Sci Technol* 2017; 54(13): 4362-4369. doi:10.1007/s13197-017-2907-y
34. Luković J, Malenčić D, Zorić L, *et al.* Anatomical characteristics and antioxidant properties of *Euphorbia nicaeensis* ssp. *glareosa*. *Open Life Sci* 2009; 4(2): 214-223. doi:10.2478/s11535-009-0003-7
35. Akhtar N, Ihsan-ul-Haq, Mirza B. Phytochemical analysis and comprehensive evaluation of antimicrobial and antioxidant properties of 61 medicinal plant species. *Arab J Chem* 2018; 11(8): 1223-1235. doi:10.1016/j.arabjc.2015.01.013
36. Battah B, Rajab A, Shbibe L, *et al.* Evaluation of antibiofilm activity of *Thymus syriacus* essential oil against clinically isolated MDR bacteria. *Prog Microbes Mol Biol* 2022; 5(1).
37. Kong GYE, Letchumanan V, Tan LTH, *et al.* Gut Microbiome in Obsessive Compulsive Disorder: Potential of Probiotics as an Adjuvant Therapy. *Prog Microbes Mol Biol* 2022; 5(1).
38. Thye AYK, Letchumanan V, Tan LTH, *et al.* Malaysia's Breakthrough in Modern Actinobacteria (MOD-ACTINO) Drug Discovery Research. *Prog Microbes Mol Biol* 2022; 5(1).
39. El Hachlafi N, Lakhdar F, Khouchlaa A, *et al.* Health Benefits and Pharmacological Properties of *Hinokitiol*. *Processes* 2021; 9(9): 1680. doi:10.3390/pr9091680
40. Bouyahya A, El Hachlafi N, Aanniz T, *et al.* Natural bioactive compounds targeting histone deacetylases in human cancers: recent updates. *Molecules* 2022; 27(8): 2568.
41. El Hachlafi N, Benkhaira N, Ferioun M, *et al.* Moroccan Medicinal Plants Used to Treat Cancer: Ethnomedicinal Study and Insights into Pharmacological Evidence. *Evid Based Complement Altern Med* 2022; 2022: 1-19. doi:10.1155/2022/1645265
42. Deng J, Xu Z, Xiang C, *et al.* Comparative evaluation of maceration and ultrasonic-assisted extraction of phenolic compounds from fresh olives. *Ultrasonics Sonochem* 2017; 37: 328-334. doi:10.1016/j.ultsonch.2017.01.023

43. Ser HL, Yin WF, Chan KG, *et al.* Antioxidant and cytotoxic potentials of *Streptomyces gilvigriseus* MUSC 26T isolated from mangrove soil in Malaysia. *Prog Microbes Mol Biol* 2018; 1(1).
44. Farhan H, Rammal H, Hijazi A, *et al.* Chemical composition and antioxidant activity of a Lebanese plant *Euphorbia macroclada schyzoceras*. *Asian Pac J Trop Biomed* 2013; 3(7): 542-548. doi:10.1016/S2221-1691(13)60110-0
45. Granato D, Shahidi F, Wrolstad R, *et al.* Antioxidant activity, total phenolics and flavonoids contents: Should we ban in vitro screening methods? *Food Chem* 2018; 264: 471-475. doi:10.1016/j.foodchem.2018.04.012
46. Janmeda PCP. Quantification of phytochemicals and in vitro antioxidant activities from various parts of *Euphorbia neriifolia* Linn. *J Appl Biol Biotechnol* 2022; 10(2): 133-145.
47. Nur Syukriah AR, Liza MS, Harisun Y, *et al.* Effect of solvent extraction on antioxidant and antibacterial activities from *Quercus infectoria* (Manjakani). *Int Food Res J* 2014; 21(3). Accessed October 19, 2022. <http://mymedr.afpm.org.my/publications/46579>
48. Yee S, Lim YC, Goh CF, *et al.* Efficacy of chlorine dioxide as a disinfectant. *Prog Microbes Mol Biol* 2020; 3(1).
49. Selvaraja M, Abdullah M, Shah AM, *et al.* Systemic Lupus Erythematosus (SLE): A review on the prevalence, clinical manifestation, and disease assessment. *Prog Microbes Mol Biol* 2020; 3(1).
50. Pham TH, Kim MS, Le MQ, *et al.* Fargesin exerts anti-inflammatory effects in THP-1 monocytes by suppressing PKC-dependent AP-1 and NF- κ B signaling. *Phytomedicine* 2017; 24: 96-103. doi:10.1016/j.phymed.2016.11.014
51. Picardo M. Isolation of Flavonoids and Flavonoid Glycosides from *Myrsine africana* and Their Inhibitory Activities against Mushroom Tyrosinase. *J Nat Prod* Published online January 4, 2018. Accessed March 6, 2023. https://www.academia.edu/50149679/Isolation_of_Flavonoids_and_Flavonoid_Glycosides_from_Myrsine_africana_and_Their_Inhibitory_Activities_against_Mushroom_Tyrosinase
52. Lahmadi S, Belhamra M, Karoune S, *et al.* Phenolic constituents and antioxidant activity of *Euphorbia retusa* Forssk. *Nat Prod Res* 2020; 34(24): 3545-3547. doi:10.1080/14786419.2019.1582040
53. Basma AA, Zakaria Z, Latha LY, *et al.* Antioxidant activity and phytochemical screening of the methanol extracts of *Euphorbia hirta* L. *Asian Pac J Trop Med* 2011; 4(5): 386-390. doi:10.1016/S1995-7645(11)60109-0
54. Mahomoodally MF, Dall'Acqua S, Sinan KI, *et al.* Phenolic compounds analysis of three *Euphorbia* species by LC-DAD-MSn and their biological properties. *J Pharm Biomed* 2020; 189: 113477. doi:10.1016/j.jpba.2020.113477
55. Yap HM, and Lye KL. An insight of vitamin E as neuroprotective agents. *Prog Microbes Mol Biol* 2020; 3(1).
56. Lee HT, Loh HC, Ramlee SNL, *et al.* Oral Dietary Supplements Use Among Healthcare Workers During the COVID-19 Pandemic in Malaysia. *Prog Microbes Mol Biol* 2021; 4(1).
57. Nanda PK, Lochan Nayak P, and Krishna Rao K. Thermal degradation analysis of biodegradable plastics from urea-modified soy protein isolate. *Polym Plast Technol Eng* 2007; 46(3): 207-211.
58. Das K, and Roychoudhury A. Reactive oxygen species (ROS) and response of antioxidants as ROS-scavengers during environmental stress in plants. *Front Environ Sci* 2014; 2: 53.
59. Cirkovic Velickovic TD, and Stanic-Vucinic DJ. The role of dietary phenolic compounds in protein digestion and processing technologies to improve their antinutritive properties. *Compr Rev Food Sci* 2018; 17(1): 82-103.